




Contents lists available at ScienceDirect

Fungal Genetics and Biology

journal homepage: www.elsevier.com/locate/yfgbiOverview of genomic and bioinformatic resources for *Zymoseptoria tritici*Alison Testa^a, Richard Oliver^a, James Hane^{a,b,*}^a Centre for Crop and Disease Management, Curtin University, Perth, WA, Australia^b Curtin Institute for Computation, Curtin University, Perth, WA, Australia

ARTICLE INFO

Article history:

Received 5 January 2015

Revised 10 April 2015

Accepted 13 April 2015

Keywords:

Zymoseptoria tritici

Genome

Transcriptome

Bioinformatics

Pathogenomics

ABSTRACT

Zymoseptoria tritici (syn. *Mycosphaerella graminicola*, *Septoria tritici*) is a haploid fungus belonging to the class Dothideomycetes. It is the causal agent of septoria leaf blotch – one of the world's most significant diseases of wheat. Here we review the genomic and bioinformatic resources that have been generated for *Z. tritici*. These include the whole-genome reference assembly for isolate IPO323, genome resequencing of alternate isolates, mitochondrial genome sequences, transcriptome sequences and expression data, and annotations of gene structure and function. We also highlight important advances in our fundamental knowledge of genome evolution and its effects on adaptation and pathogenicity in *Z. tritici* that have been facilitated by these resources.

© 2015 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. The nuclear genome

An early landmark paper studied the electrophoretic karyotype of 7 isolates of *Z. tritici*, which estimated 14–16 chromosomes ranging from 330 kb to 3.5 Mb (McDonald and Martinez, 1991). Chromosome length and number polymorphisms were also observed between isolates, with chromosome presence/absence variation (PAV) observed exclusively for the two shortest chromosome bands. Subsequent generation of a genetic map from a cross between parent isolates IPO323 and IPO94269 predicted a higher number of chromosomes at 23 linkage groups (Kema et al., 2002). Further analysis of this mapping population revealed that at least 8 of the smaller linkage groups – corresponding to 8 dispensable chromosomes ranging from 0.39 to 0.77 Mb – were not required for saprophytic growth i.e. were dispensable or accessory chromosomes (ACs) (Wittenberg et al., 2009). The distinction between core chromosomes (CCs) and ACs was an important milestone for *Z. tritici* genomics.

Z. tritici was the first species of the Dothideomycetes – a fungal class of high agricultural significance (Hane et al., 2011b; Ohm et al., 2012) – to have had the genome of a representative isolate (IPO323) (Kema and van Silfhout, 1997) sequenced beyond the typical 'draft genome' status, achieving near complete

chromosome sequences spanning from telomere to telomere (Goodwin et al., 2011) (Table 1). The IPO323 reference genome assembly is 39.69 Mb in length, representing 21 chromosomes. At least half of the genome is contained within the six largest assembled sequences (that is, an N50 of 6), with the sixth largest sequence having a length of 2.67 Mb. Its chromosomes have been designated into CCs and ACs, numbered 1–13 and 14–21 respectively.

In addition to chromosomal polymorphisms across the ACs, the *Z. tritici* genome has also been observed to exhibit at least two other forms of genomic plasticity. The repetitive content of the IPO323 genome reference is 12.26% (Ohm et al., 2012) and its repeats exhibit the hallmarks of repeat-induced point mutation (RIP) (Goodwin et al., 2011) – a fungal specific type of mutation that targets repetitive DNA and randomly converts cytosine to thymine bases (Hane and Oliver, 2008, 2010; Hane et al., 2015). In comparisons of whole-genome sequences across species belonging to the sub-phylum Pezizomycotina, the *Z. tritici* genome also exhibits a typical "mesosyntentic" conservation pattern, that is, the relative order and orientation of genes are reshuffled within homologous chromosomes due to frequent intra-chromosomal recombinations (Croll et al., 2013; Hane et al., 2011a). The combined effects of AC variability, mesosynteny and RIP may have significant implications for genome evolution and pathogenicity in *Z. tritici*.

Due to the completeness of its genome assembly, *Z. tritici* IPO323 has been a key point of reference in several comparative genomics studies, particularly across the Dothideomycetes (Hane et al., 2011b; Morais do Amaral et al., 2012; Ohm et al., 2012). Observations of mesosynteny made from genome sequence

* Corresponding author at: CCDM Bioinformatics, Centre for Crop and Disease Management, Department of Environment and Agriculture, Curtin University, Australia. Tel.: +61 8 9266 1726. Postal address: Department of Environment & Agriculture, Curtin University, GPO Box U1987, Perth, Western Australia 6845, Australia.

E-mail address: James.Hane@curtin.edu.au (J. Hane).

alignments of different fungal species with *Z. tritici* can also be applied to fungal genome finishing. Whole-genome alignments of draft assemblies of other Pezizomycotina species with the *Z. tritici* IPO323 reference assembly have previously been used to predict the co-location of scaffolds on the same chromosome (i.e. synteny) (Goodwin et al., 2011). Significant insight has also been gleaned within the species *Z. tritici* itself through comparisons of alternate isolates, revealing differential selection pressures between CCs and ACs (Stukenbrock et al., 2010) and highlighting isolate-specific patterns of chromosome presence and absence (Goodwin et al., 2011; Croll et al., 2013) (see also McDonald et al., 1991).

2. The mitochondrial genome

The mitochondrial genomes (mtDNA) of two *Z. tritici* isolates, the reference isolate IPO323 and strain STBB1 (Table 1), have been sequenced in full (Torriani et al., 2008). The *Z. tritici* mtDNA [Nucleotide: EU090238] is 43,960 bp in length, indicating that it has been relatively untouched by invading intronic endonucleases (as compared with *Leptosphaeria maculans* and *Pyrenophora tritici-repentis*) (Manning et al., 2013; Rouxel et al., 2011). The mtDNA contains 15 protein-coding genes, large and small ribosomal subunits, 27 tRNAs and 8 unknown open-reading frames. Protein-coding genes residing on the mtDNA include *atp6*, *atp8*, *atp9*, *cox1-3*, *cytb*, *nad1-6*, *nad4L* and *RNA-Pol*. The presence and relative order of these genes is not highly conserved with closely related species of the Dothideomycetes (Aguileta et al., 2014; Hane et al., 2011b). However within the species *Z. tritici* across multiple isolates there was relatively little sequence variability (Zhan et al., 2003). Polymorphic sequences were identified between the mtDNA of the two sequenced isolates and used to generate markers for PCR screening across numerous isolates – an important genomic resource for studying the emergence and distribution of fungicide resistance. The mtDNA sequence data has also been a useful tool for tracking the history of the co-evolution of *Z. tritici* and its wheat host (Torriani et al., 2011; Zhan et al., 2004).

3. Gene function

Annotations of gene structure and function for *Z. tritici* IPO323 (Goodwin et al., 2011) are downloadable from JGI Mycocosm (Grigoriev et al., 2013) and EnsemblFungi (Kersey et al., 2014) (Table 1). As part of the initial genome survey study, potential carbohydrate-active enzymes (CAZymes) encoded by the gene content of *Z. tritici* were annotated, highlighting a distinctive reduction in cellulose and other cell-wall degrading enzymes relative to other plant pathogens (Goodwin et al., 2011). This observation supported a model of “stealth pathogenesis” for *Z. tritici*, in which CAZyme activity is reduced – with a compensatory shift toward protein degradation (Goodwin et al., 2011) – in conjunction with other adaptations for avoidance of triggering host-defenses (Lee et al., 2014). Morais do Amaral et al. subsequently used the gene annotations generated by Goodwin et al. to bioinformatically predict genes encoding secreted proteins and assign additional functional annotations. The predicted secretome (Morais do Amaral et al., 2012) comprises 492 proteins, with 321 possessing some level of functional annotation and 171 with no functional annotation.

4. Gene expression

An early transcriptome resource for *Z. tritici* was generated by Kema et al., comprising Expressed Sequence Tag (EST) libraries for IPO323 from 7 *in vitro* and 3 *in planta* growth conditions (Kema et al., 2008) (Table 1). Limitations around capture of fungal

transcripts in mixed plant samples were overcome by purifying via hybridisation against IPO323 genomic DNA. A total of 27,007 ESTs were clustered into 9190 unigene loci, totaling 5.2 Mb. Though a comparable number of ESTs were typically sequenced in similar fungal genomics studies of the day, a very high representation of loci was achieved due to the high number of libraries from diverse growth conditions.

Recently, high-coverage whole-transcriptome expression data for the *Z. tritici*-wheat interaction has been facilitated by next-generation sequencing (RNA-Seq) (Table 1). Yang et al. have generated 18 Gbp of raw RNA-Seq data, comparing expression across the infection and necrotrophic growth stages of *Z. tritici* at 4, 10 and 13 days post-infection (dpi), as well as the corresponding host responses in *Triticum aestivum* [BioProject: 196595] (Yang et al., 2013). This study expanded upon the predicted secretome of Morais do Amaral et al., highlighting 313 genes of IPO323 that were highly expressed during early and late infection. A second RNA-Seq study by Kellner et al., generating 117.8 Gbp of RNA-Seq data, followed up with an in-depth comparison of *Z. tritici* expression during infection of two hosts: *T. aestivum* and *Brachypodium distachyon* [NCBI GEO: GSE54874] (Kellner et al., 2014). Differential expression patterns were observed across core and accessory chromosomes of *Z. tritici*. Kellner et al. observed no host-specific gene expression on accessory chromosomes during early infection and 25 wheat-specific genes during late infection (13 dpi). Notably, neither study had used the wealth of exon structure information afforded by RNA-Seq to re-annotate the initial set of gene predictions presented by Goodwin et al. (2011) (which relied heavily on *in silico* gene predictions supplemented with limited EST and homology supporting data). Consequently, despite considerable accumulation of knowledge of conditional gene-expression, a reasonable level of error in the prediction of gene loci, their exon structure and their translated sequences has persisted in the gene-based datasets of *Z. tritici* for a few years. However, RNA-seq data generated in a recent transcriptome study (Rudd et al., 2015) has been used in-house by the same research group to improve the accuracy of gene annotations in *Z. tritici* (annotations available from EnsemblFungi) (Kersey et al., 2014). The study itself by Rudd et al. has provided further insight into the wheat-*Z. tritici* interaction. RNA-Seq was used to profile expression at five infection time points (symptomless growth: 1 and 4 dpi; host-cell death: 9 and 14 dpi; and asexual sporulation: 21 dpi), detecting expression of at least 80% of predicted loci and identifying >3000 *Z. tritici* genes and >7000 wheat genes as differentially expressed during infection. Variation in the contribution of chromosomes to changes in expression was observed, with CCs exhibiting the highest overall and most significant changes in expression. Conversely ACs exhibited minimal gene expression and few differentially expressed genes. Genes encoding candidate effector proteins were found to be up-regulated *in planta* and expression data was used provide additional supporting evidence for the “top” candidates. Incorporating the above with profiles of differential metabolites over the same time course, Rudd et al. also observed that the switch from latent to necrotrophic growth and reproduction coincided with the activation of plant defense responses and a switch in *Z. tritici*'s nutrient source from lipids and fatty acid stores to carbohydrate metabolism.

5. Genome mutation, adaptation and pathogenicity

ACs have been described by Croll and McDonald as a “cradle for adaptive evolution”. This refers to their potential for retention of higher levels of mutation over time, due to low impact on fitness and correspondingly lower selective pressures (Croll and McDonald, 2012). A two-speed rate of evolution across CCs and




Fig. 1. Histograms of the GC content of segmented regions of the *Z. tritici* IPO323 genome assembly, and the relative proportion of the genome vs GC content. (A) The dotted lines show the two components of the Cauchy mixture model fit to the data. The solid vertical line shows the GC content selected to categorize segments, corresponding to the intersection between the two components of the mixture model. (B) Relative proportions of the *Z. tritici* IPO323 CC and AC chromosome subsets corresponding to ranges of GC content from 35% to 65% (red), illustrating depleted GC content within accessory chromosomes. The solid vertical line (blue) shows the GC content selected to categorize segmented regions of the genome as AT-rich or GC-equilibrated.

ACs allows pathogens like *Z. tritici* to retain its core gene content governing its viability and metabolism, whilst at the same time innovating and mutating genes rapidly in response to host-defences and control strategies. Plant pathogenicity genes have previously been observed to reside on the ACs of other plant-pathogenic fungi species, including *F. oxysporum* f. sp. *lycopersici* (Ma et al., 2010), *F. solani* (Coleman et al., 2009). Despite this, no *Z. tritici* pathogenicity genes have been mapped to ACs (Goodwin et al., 2011) with recent work reporting eight QTLs, all of which mapped to CCs (Mirzadi Gohari et al., 2015).

Z. tritici ACs are richer in repetitive DNA relative to CCs and also undergo recombination (including mesosyntentic rearrangements) more frequently (Croll et al., 2013). The higher repetitive DNA content of ACs, combined with active RIP, can also lead to the rapid mutation of non-repetitive genes flanking RIP-targeted repeats. In the related Dothideomycete plant pathogen *Leptosphaeria maculans* – which is well known for widespread distribution of AT-rich regions (syn. AT-rich isochores) within its genome – the “leakage” of RIP into neighboring non-repetitive has been demonstrated to accelerate the rate of non-synonymous mutation in avirulence genes arranged in a repeat-proximal configuration (Fudal et al., 2009; Hane et al., 2015; Van de Wouw et al., 2010). The *Z. tritici* genome has a similar abundance of AT-rich regions across its genome, particularly within its ACs, which may also have contributed to gene innovation and diversification over time. To assess

Table 1

Summary of sequence and bioinformatic resources available for *Z. tritici*.

Resource	Repository	Reference
EST libraries (IPO323)	EMBL (multiple accessions)	Kema et al. (2008)
Nuclear genome (IPO323)	NCBI Nucleotide: ACPE000000000.1	Goodwin et al. (2011)
Mitochondrial genome (IPO323 & STBB1)	NCBI Nucleotide: EU090238	Torriani et al. (2011, 2008)
Gene annotation (IPO323)	JGI (www.jgi.doe.gov)	Goodwin et al. (2011), Grigoriev et al. (2013)
Predicted secretome (IPO323)	EnsemblFungi n/a	Kersey et al. (2014) Morais do Amaral et al. (2012)
Gene expression: RNA-Seq data for host and pathogen during wheat infection	NCBI BioProject: 196595 NCBI BioProject: 278138	Rudd et al. (2015)
Gene expression: RNA-Seq data for wheat infection vs <i>B. distachyon</i> infection	NCBI GEO: GSE54874	Kellner et al. (2014)
Mapped QTLs and effector candidate list (IPO323)	n/a	Mizardi Gohari et al. (2015)
Resequencing	n/a	Croll et al. (2013), McDonald et al. (1991)

the prevalence of AT-rich regions in *Z. tritici*, the genome was recursively segmented into regions of differing GC content using Jensen-Shannon divergence (Bernaola-Galván et al., 1996; Elhaik et al., 2010a, 2010b) (stopping criteria: minimum segment length of 1000 bp, t-test (5% significance level) on adjacent average GC values, similar to Oliver et al. (2004)). Segmented genome regions were classified according to their constituent GC content. A mixture of two Cauchy distributions was fit to the data using expectation maximization, which identified two peaks centered at 54.2% and 44.2% GC. This allowed a GC boundary to be defined at 49.3% GC in order to distinguish “GC-equilibrated” from “AT-rich” DNA regions (Fig. 1A, Supplementary Data File 1). The AT-rich regions are substantially shorter than GC-equilibrated regions, with average lengths of 10.8 kbp and 48.5 kbp respectively. AT-rich regions were observed on all chromosomes and comprised 18.2% of the total genome length, however the proportion differed between ACs to CCs with AT-rich regions comprising 31.9% in ACs and 16.3% in CCs (Fig. 1B). The majority of annotated genes reside within GC-equilibrated regions, with an overall gene density of 334 genes/Mbp. In contrast, 91 genes have $\geq 50\%$ of their length within AT-rich regions and are comparatively sparsely distributed at 12.6 genes/Mbp. Studies of related Dothideomycete plant pathogens, e.g. *Leptosphaeria maculans* and *Passalora fulva*, contain many secreted and/or effector-like genes associated with similar AT-rich regions (de Wit et al., 2012; Rouxel et al., 2011; Van de Wouw et al., 2010). As such, although genes within AT-rich regions contribute little to the overall gene content or gene-expression of *Z. tritici*, they may still play some role in pathogenicity and adaptability.

6. Conclusion

The sum of sequence and bioinformatic resources generated over the last decade for *Z. tritici* (Table 1) has significantly advanced knowledge of this fungal pathogen and its host-interactions. Foremost among these has been the generation of the whole-genome sequence – a solid foundation upon which multiple layers of additional information have been juxtaposed.

Various bioinformatic techniques have been applied to predicted gene regions and their protein products with a view to focussing on those with properties relevant to pathogenicity. This has been complemented by analysis of gene expression during infection, which has highlighted generic and host-specific pathogenicity genes in *Z. tritici*. The chromosome-length genome sequence has itself been a powerful tool in understanding genome mutation mechanisms influencing adaptation in *Z. tritici* and assessing the “genomic context” of its putative pathogenicity genes.

Appendix A. Supplementary material

Supplementary data associated with this article can be found, in the online version, at <http://dx.doi.org/10.1016/j.fgb.2015.04.011>.

References

- Aguileta, G. et al., 2014. High variability of mitochondrial gene order among fungi. *Genome Biol. Evol.* 6, 451–465.
- Bernaola-Galván, P. et al., 1996. Compositional segmentation and long-range fractal correlations in DNA sequences. *Phys. Rev. E* 53, 5181.
- Coleman, J.J. et al., 2009. The genome of *Nectria haematococca*: contribution of supernumerary chromosomes to gene expansion. *PLoS Genet.* 5, e1000618.
- Croll, D., McDonald, B.A., 2012. The accessory genome as a cradle for adaptive evolution in pathogens. *PLoS Pathog.* 8, e1002608.
- Croll, D. et al., 2013. Breakage-fusion-bridge cycles and large insertions contribute to the rapid evolution of accessory chromosomes in a fungal pathogen. *PLoS Genet.* 9, e1003567.
- de Wit, P.J. et al., 2012. The genomes of the fungal plant pathogens *Cladosporium fulvum* and *Dothistroma septosporium* reveal adaptation to different hosts and lifestyles but also signatures of common ancestry. *PLoS Genet.* 8, e1003088.
- Elhaik, E. et al., 2010a. Comparative testing of DNA segmentation algorithms using benchmark simulations. *Mol. Biol. Evol.* 27, 1015–1024.
- Elhaik, E. et al., 2010b. Identifying compositionally homogeneous and nonhomogeneous domains within the human genome using a novel segmentation algorithm. *Nucleic Acids Res.* 38, e158–e158.
- Fudal, I. et al., 2009. Repeat-induced point mutation (RIP) as an alternative mechanism of evolution toward virulence in *Leptosphaeria maculans*. *Mol. Plant Microbe Interact.* 22, 932–941.
- Goodwin, S.B. et al., 2011. Finished genome of the fungal wheat pathogen *Mycosphaerella graminicola* reveals dispensome structure, chromosome plasticity, and stealth pathogenesis. *PLoS Genet.* 7, e1002070.
- Grigoriev, I.V. et al., 2013. MycoCosm portal: gearing up for 1000 fungal genomes. *Nucl. Acids Res.*, gkt1183.
- Hane, J.K., Oliver, R.P., 2008. RIPCAL: a tool for alignment-based analysis of repeat-induced point mutations in fungal genomic sequences. *BMC Bioinform.* 9, 478.
- Hane, J.K., Oliver, R.P., 2010. *In silico* reversal of repeat-induced point mutation (RIP) identifies the origins of repeat families and uncovers obscured duplicated genes. *BMC Genom.* 11, 655.
- Hane, J.K. et al., 2011a. A novel mode of chromosomal evolution peculiar to filamentous Ascomycete fungi. *Genome Biol.* 12, R45.
- Hane, J.K. et al., 2011b. Genomic and comparative analysis of the class dothideomycetes. *Evolution of Fungi and Fungal-Like Organisms*. Springer, pp. 205–229.
- Hane, J.K. et al., 2015. Repeat-Induced Point Mutation: A Fungal-Specific, Endogenous Mutagenesis Process. *Genetic Transformation Systems in Fungi*, vol. 2. Springer, pp. 55–68.
- Kellner, R. et al., 2014. Expression profiling of the wheat pathogen *Zymoseptoria tritici* reveals genomic patterns of transcription and host-specific regulatory programs. *Genom. Biol. Evol.* 6, 1353–1365.
- Kema, G.H., van Silfhout, C.H., 1997. Genetic variation for virulence and resistance in the Wheat-*Mycosphaerella graminicola* Pathosystem III. Comparative seedling and adult plant experiments. *Phytopathology* 87, 266–272.
- Kema, G.H. et al., 2002. A combined amplified fragment length polymorphism and randomly amplified polymorphism DNA genetic linkage map of *Mycosphaerella graminicola*, the septoria tritici leaf blotch pathogen of wheat. *Genetics* 161, 1497–1505.
- Kema, G.H. et al., 2008. Large-scale gene discovery in the septoria tritici blotch fungus *Mycosphaerella graminicola* with a focus on *in planta* expression. *Mol. Plant Microbe Interact.* 21, 1249–1260.
- Kersey, P.J. et al., 2014. Ensembl Genomes 2013: scaling up access to genome-wide data. *Nucleic Acids Res.* 42, D546–D552.
- Lee, W.S. et al., 2014. *Mycosphaerella graminicola* LysM effector-mediated stealth pathogenesis subverts recognition through both CERK1 and CEBIP homologues in wheat. *Mol. Plant Microbe Interact.* 27, 236–243.
- Ma, L.J. et al., 2010. Comparative genomics reveals mobile pathogenicity chromosomes in *Fusarium*. *Nature* 464, 367–373.
- Manning, V.A. et al., 2013. Comparative genomics of a plant-pathogenic fungus, *Pyrenophora tritici-repentis*, reveals transduplication and the impact of repeat elements on pathogenicity and population divergence. *G3: Genes| Genomes| Genetics* 3, 41–63.
- McDonald, B., Martinez, J., 1991. Chromosome length polymorphisms in a *Septoria tritici* population. *Curr. Genet.* 19, 265–271.
- Mirzadi Gohari, A., 2015. Effector discovery in the fungal wheat pathogen *Zymoseptoria tritici*. *Mol. Plant Pathol.* <http://dx.doi.org/10.1111/mpp.12251>.
- Morais do Amaral, A. et al., 2012. Defining the predicted protein secretome of the fungal wheat leaf pathogen *Mycosphaerella graminicola*. *PLoS ONE* 7, e49904.
- Ohm, R.A. et al., 2012. Diverse lifestyles and strategies of plant pathogenesis encoded in the genomes of eighteen Dothideomycetes fungi. *PLoS Pathog.* 8, e1003037.
- Oliver, J.L. et al., 2004. IsoFinder: computational prediction of isochores in genome sequences. *Nucleic Acids Res.* 32, W287–W292.
- Rouxel, T. et al., 2011. Effector diversification within compartments of the *Leptosphaeria maculans* genome affected by Repeat-Induced Point mutations. *Nature Commun.* 2, 202.
- Rudd, J. et al., 2015. Transcriptome and metabolite profiling the infection cycle of *Zymoseptoria tritici* on wheat (*Triticum aestivum*) reveals a biphasic interaction with plant immunity involving differential pathogen chromosomal contributions, and a variation on the hemibiotrophic lifestyle definition. *Plant Physiol.*, pp. 114.255927.
- Stukenbrock, E.H. et al., 2010. Whole-genome and chromosome evolution associated with host adaptation and speciation of the wheat pathogen *Mycosphaerella graminicola*. *PLoS Genet.* 6, e1001189.
- Torriani, S.F. et al., 2008. Intraspecific comparison and annotation of two complete mitochondrial genome sequences from the plant pathogenic fungus *Mycosphaerella graminicola*. *Fungal Genet. Biol.* 45, 628–637.
- Torriani, S.F. et al., 2011. Evolutionary history of the mitochondrial genome in *Mycosphaerella* populations infecting bread wheat, durum wheat and wild grasses. *Mol. Phylogenet. Evol.* 58, 192–197.
- Van de Wouw, A.P. et al., 2010. Evolution of linked avirulence effectors in *Leptosphaeria maculans* is affected by genomic environment and exposure to resistance genes in host plants. *PLoS Pathog.* 6, e1001180.
- Wittenberg, A.H. et al., 2009. Meiosis drives extraordinary genome plasticity in the haploid fungal plant pathogen *Mycosphaerella graminicola*. *PLoS ONE* 4, e5863.
- Yang, F. et al., 2013. Transcriptional reprogramming of wheat and the hemibiotrophic pathogen *Septoria tritici* during two phases of the compatible interaction. *PLoS ONE* 8, e81606.
- Zhan, J. et al., 2003. The global genetic structure of the wheat pathogen *Mycosphaerella graminicola* is characterized by high nuclear diversity, low mitochondrial diversity, regular recombination, and gene flow. *Fungal Genet. Biol.* 38, 286–297.
- Zhan, J. et al., 2004. Evidence for natural selection in the mitochondrial genome of *Mycosphaerella graminicola*. *Phytopathology* 94, 261–267.